



ARSET Applied Remote Sensing Training http://arset.gsfc.nasa.gov

@NASAARSET

Smoke Monitoring From Space

Pawan Gupta and Melanie Follette-Cook

Satellite Remote Sensing of Air Quality

September 19-21, 2017 University of California, Riverside


www.nasa.gov

Learning Objectives

By the end of this presentation, you will be able to:

- describe existing satellite capabilities for smoke monitoring
- describe available smoke products and their applications

Fires in Pictures – Google Image Search



Forest Fires in Pictures - Google Image Search




Agriculture Fires in Pictures - Google Image Search



Waste Burning in Pictures - Google Image Search




Importance of Smoke and Fire Monitoring




Fire Detection From Satellites


- By detecting smoke
- By detecting temperature anomaly
- By detecting light









Visible Smoke From Fires



Spectral Signatures - Smoke Over Land




Clear sky Smoke Heavy smoke Clouds




Zhao et al., 2010

Spectral Signatures - Smoke Over Ocean




Clear sky Smoke Heavy smoke Clouds




Zhao et al., 2010

Smoke Detection Example (Zhao et al., 2010)




Smoke Monitoring Tools – Worldview



- Visible Imagery (MODIS, VIIRS)
- Fire Detection (MODIS, VIIRS)
- Aerosol Optical Depth (MODIS, OMI, MISR)
- Aerosol Index (OMI)
- Day-Night Band (VIIRS)

Smoke Monitoring Tools – eIDEA



- Visible Imagery (VIIRS, only US & Canada)
- Smoke Mask
- Aerosol Optical Depth
- AirNOW PM2.5
- Satellite derived PM2.5


NOAA's Hazard Mapping System




http://www.ospo.noaa.gov/Products/land/hms.html

Smoke Monitoring Tools – MISR Plume Height

https://misr.jpl.nasa.gov/getData/accessData/MisrMinxPlumes2/



GOES-R or GOES-16



Other Tools

NASA's GEOS-5 Aerosol Forecasts

https://portal.nccs.nasa.gov/cgi-fp/fp_2d_chem.cgi

NRL Forecasts


https://www.nrlmry.navy.mil/aerosol/#currentaerosolmodeling



Hyperion Hyperspectral Imagery Showing Tucson Wildfires - 3 July 2003







$$BI = \frac{\rho_{1100} - \rho_{2200}}{\rho_{1100} + \rho_{2200}}$$

The Burn Index (BI) is used to detect burn scars due to wildfires. It has the Advantage over the NDVI in that it uses wavelengths that are transparent to smoke (1.1 and 2.2 microns).

Change is Reflectance After Fire

